

## African Plants

[Red List statistics](#)  
[methodology](#)
[Summary of recent changes](#)  
[Glossary](#)  
[Guidelines for EIAs](#)
[National Red List categories](#)
[Assessment](#)

### Browse

- [Genera: A](#)
- [Genera: B](#)
- [Genera: C](#)
- [Genera: D](#)
- [Genera: E](#)
- [Genera: F](#)
- [Genera: G](#)
- [Genera: H](#)
- [Genera: I](#)
- [Genera: J](#)
- [Genera: K](#)
- [Genera: L](#)
- [Genera: M](#)
- [Genera: N](#)
- [Genera: O](#)
- [Genera: P](#)
- [Genera: Q](#)
- [Genera: R](#)
- [Genera: S](#)
- [Genera: T](#)
- [Genera: U](#)
- [Genera: V](#)
- [Genera: W](#)
- [Genera: X](#)
- [Genera: Y](#)
- [Genera: Z](#)

[Home](#) >> [Genera: T](#) >> [Genus: Tylecodon](#)

### Taxonomy

Scientific Name	<b>Tylecodon hallii</b> (Toelken) <b>Toelken</b>
Higher Classification	Dicotyledons
Family	CRASSULACEAE
Synonyms	Cotyledon hallii Toelken, Cotyledon pearsonii Schönland (in part), Cotyledon racemosa E.Mey. ex Harv. (in part)



### National Status

Status and Criteria	<b>Least Concern</b>
Assessment Date	2015/05/12
Assessor(s)	L. von Staden
Justification	A range-restricted, but locally common species that is not in danger of extinction.



© A.G. Rebelo



### Distribution

Endemism	Not endemic to South Africa
Provincial distribution	Northern Cape
Range	Southern Namibia to the eastern Richtersveld.

 Search for images of **Tylecodon hallii** on [iNaturalist](#)

### Habitat and Ecology

Major system	Terrestrial
Major habitats	Desert, Succulent Karoo
Description	Exposed rocky slopes.

## Population

This species is common across the eastern Richtersveld between Sendelingsdrif and Steinkopf. In Namibia it is known from 10 to 20 subpopulations, and not declining (Loots 2005).

**Population** Stable trend

## Assessment History

Taxon assessed and Criteria	Status	Citation/Red List version
Tylecodon hallii (Toelken)	<b>Least Concern</b>	Raimondo et al. (2009)

## Bibliography

Loots, S. 2005. Red Data Book of Namibian plants. Southern African Botanical Diversity Network Report 38. SABONET, Pretoria and Windhoek.

Raimondo, D., von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A. and Manyama, P.A. 2009. Red List of South African Plants. Strelitzia 25. South African National Biodiversity Institute, Pretoria.

Snijman, D.A. 2013. Plants of the Greater Cape Floristic Region 2: The extra Cape flora. Strelitzia 30. South African National Biodiversity Institute, Pretoria.

Tölken, H.R. 1977. New taxa and a new combination in the genus *Cotyledon*. Bothalia 12(2):191-194.

Tölken, H.R. 1978. New taxa and new combinations in *Cotyledon* and allied genera. Bothalia 12(3):377-393.

Tölken, H.R. 1985. Crassulaceae. In: O.A. Leistner (ed). Flora of southern Africa 14, 1:1-244. Botanical Research Institute, Pretoria.

Van Jaarsveld, E.J. and Koutnik, D. 2004. *Cotyledon* and *Tylecodon*. Umdaus Press, Pretoria.

## Citation

---

von Staden, L. 2015. *Tylecodon hallii* (Toelken) Toelken. National Assessment: Red List of South African Plants version 2020.1. Accessed on 2021/12/01

 [Comment on this assessment](#)

(c) South African National Biodiversity Institute ([SANBI](#)) 2010-12.