

Red List of South

[Search the Red List](#)

African Plants

[Red List statistics](#)
[methodology](#)
[Summary of recent changes](#)
[Glossary](#)
[Guidelines for EIAs](#)
[National Red List categories](#)
[Assessment](#)
Browse
[Home >> Genera: C >> Genus: Conophytum](#)

- [Genera: A](#)
- [Genera: B](#)
- [Genera: C](#)
- [Genera: D](#)
- [Genera: E](#)
- [Genera: F](#)
- [Genera: G](#)
- [Genera: H](#)
- [Genera: I](#)
- [Genera: J](#)
- [Genera: K](#)
- [Genera: L](#)
- [Genera: M](#)
- [Genera: N](#)
- [Genera: O](#)
- [Genera: P](#)
- [Genera: Q](#)
- [Genera: R](#)
- [Genera: S](#)
- [Genera: T](#)
- [Genera: U](#)
- [Genera: V](#)
- [Genera: W](#)
- [Genera: X](#)
- [Genera: Y](#)
- [Genera: Z](#)

Taxonomy

Scientific Name	Conophytum uviforme (Haw.) N.E.Br. subsp. uviforme
Higher Classification	Dicotyledons
Family	AIZOACEAE
Synonyms	Conophytum clarum N.E.Br., Conophytum colorans Lavis, Conophytum framesii Lavis, Conophytum francisci L.Bolus, Conophytum hillii L.Bolus, Conophytum julii Schwantes ex H.Jacobsen, Conophytum litorale L.Bolus, Conophytum meleagris L.Bolus, Conophytum pardivisum Tischer, Conophytum praeparvum N.E.Br., Conophytum praeparvum N.E.Br. var. praeparvum, Conophytum praeparvum N.E.Br. var. roseum Lavis, Conophytum prolongatum L.Bolus, Conophytum rooipanense L.Bolus, Conophytum stipitatum L.Bolus, Conophytum uviforme (Haw.) N.E.Br. forma framesii (Lavis) Tischer, Conophytum uviforme (Haw.) N.E.Br. forma meleagris (L.Bolus) Tischer, Conophytum uviforme (Haw.) N.E.Br. var. clarum (N.E.Br.) Rawé, Conophytum uviforme (Haw.) N.E.Br. var. litorale (L.Bolus) Rawé, Conophytum



© A.J. Young

 Search for images of **Conophytum uviforme subsp. uviforme** on [iNaturalist](#)

uviforme (Haw.)
 N.E.Br. var. occultum
 (L.Bolus) Rawé,
Conophytum uviforme
 (Haw.) N.E.Br. var.
subincanum (Tischer)
 Rawé, *Conophytum*
vanrhynsdorpense
 Schwantes,
Mesembryanthemum
uviforme (Haw.)
 N.E.Br.

National Status

Status and Criteria	Least Concern
Assessment Date	2015/06/10
Assessor(s)	A.J. Young, J.E. Victor & P.G. Desmet
Justification	This taxon is widespread, common and not in danger of extinction.

Distribution

Endemism	South African endemic
Provincial distribution	Northern Cape, Western Cape
Range	Vredendal to Calvinia and Loeriesfontein.

Habitat and Ecology

Major system	Terrestrial
Major habitats	Fynbos, Succulent Karoo
Description	Shale flats with quartz rubble, but also in crevices on gneiss domes, as well as granite and quartz outcrops.

Threats

There are no significant threats to this taxon.

Population

This taxon is widespread and known from many subpopulations. It is often locally abundant, and the population size is likely to be well in excess of 10 000 plants.

Population trend Stable

Assessment History

Taxon assessed	Status and Criteria	Citation/Red List version

<i>Conophytum</i>	Least	Raimondo et al. (2009)
uviforme	Concern	
(Haw.)		
N.E.Br.		
subsp.		
uviforme		
<i>Conophytum</i>	Vulnerable	Hilton-Taylor (1996)
uviforme		
(Haw.)		
N.E.Br. var.		
subincanum		
(Tischer)		
Rawé		
<i>Conophytum</i>	Rare	Hilton-Taylor (1996)
julii		
Schwantes		
ex		
H.Jacobsen		

Bibliography

Hammer, S. 2002. Dumpling and his wife: New view of the genus *Conophytum*. EAE Creative Colour, Norwich.

Hammer, S.A. 1993. The genus *Conophytum*: A conograph. Succulent Plant Publications, Pretoria.

Hilton-Taylor, C. 1996. Red data list of southern African plants. *Strelitzia* 4. South African National Botanical Institute, Pretoria.

Raimondo, D., von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A. and Manyama, P.A. 2009. Red List of South African Plants. *Strelitzia* 25. South African National Biodiversity Institute, Pretoria.

Citation

Young, A.J., Victor, J.E. & Desmet, P.G. 2015. *Conophytum uviforme* (Haw.) N.E.Br. subsp. *uviforme*. National Assessment: Red List of South African Plants version 2020.1. Accessed on 2021/10/11

 [Comment on this assessment](#)