


African Plants

[Red List statistics](#)
[methodology](#)
[Summary of recent changes](#)
[Glossary](#)
[Guidelines for EIAs](#)
[National Red List categories](#)
[Assessment](#)
Browse
[Home](#) >> [Genera: C](#) >> [Genus: Conophytum](#)

- [Genera: A](#)
- [Genera: B](#)
- [Genera: C](#)
- [Genera: D](#)
- [Genera: E](#)
- [Genera: F](#)
- [Genera: G](#)
- [Genera: H](#)
- [Genera: I](#)
- [Genera: J](#)
- [Genera: K](#)
- [Genera: L](#)
- [Genera: M](#)
- [Genera: N](#)
- [Genera: O](#)
- [Genera: P](#)
- [Genera: Q](#)
- [Genera: R](#)
- [Genera: S](#)
- [Genera: T](#)
- [Genera: U](#)
- [Genera: V](#)
- [Genera: W](#)
- [Genera: X](#)
- [Genera: Y](#)
- [Genera: Z](#)

Taxonomy

Scientific Name	Conophytum truncatum (Thunb.) N.E.Br. subsp. <i>viridicatum</i> (N.E.Br.) S.A.Hammer
Higher Classification	Dicotyledons
Family	AIZOACEAE
Synonyms	Conophytum archeri Lavis (in part), Conophytum archeri Lavis var. stayneri L.Bolus, Conophytum catervum (N.E.Br.) N.E.Br., Conophytum complanatum L.Bolus, Conophytum dispar N.E.Br., Conophytum koupense Tischer, Conophytum longitubum L.Bolus, Conophytum muirii N.E.Br., Conophytum novellum N.E.Br., Conophytum <i>pisinnum</i> N.E.Br., Conophytum <i>pisinnum</i> (N.E.Br.) N.E.Br. var. <i>pisinnum</i> (N.E.Br.) Rawé, Conophytum <i>viridicatum</i> (N.E.Br.) N.E.Br. var. <i>pisinnum</i> (N.E.Br.) Rawé, Conophytum <i>viridicatum</i> (N.E.Br.) N.E.Br. var. <i>punctatum</i> N.E.Br., Conophytum <i>viridicatum</i> (N.E.Br.) N.E.Br. var. <i>viridicatum</i> , Mesembryanthemum catervum N.E.Br., Mesembryanthemum <i>pisinnum</i> N.E.Br.



© A.J. Young

Search for images of **Conophytum truncatum** subsp. ***viridicatum*** on [iNaturalist](#)

National Status

Status and Criteria	Least Concern
Assessment Date	2015/06/10
Assessor(s)	A.J. Young
Justification	Widespread, common and not in danger of extinction.

Distribution

Endemism	South African endemic
Provincial distribution	Western Cape
Range	Western Little Karoo, from Montagu to Laingsburg.

Habitat and Ecology

Major system	Terrestrial
Major habitats	Succulent Karoo
Description	Quartzite scree and summits of shale ridges.

Threats

There are no significant threats to this taxon.

Population

This taxon is widespread and often locally abundant. The population is not declining.

Population trend	Stable
-------------------------	--------

Assessment History

Taxon assessed	Status and Criteria	Citation/Red List version
<i>Conophytum truncatum</i> (Thunb.) N.E.Br. subsp. <i>viridicatum</i> (N.E.Br.) S.A.Hammer	Least Concern	Raimondo et al. (2009)

Bibliography

Hammer, S. 2002. Dumpling and his wife: New view of the genus *Conophytum*. EAE Creative Colour, Norwich.

Hammer, S.A. 1993. The genus *Conophytum*: A conograph.

**Succulent Plant Publications,
Pretoria.**

Raimondo, D., von Staden, L.,
Foden, W., Victor, J.E., Helme,
N.A., Turner, R.C., Kamundi, D.A.
and Manyama, P.A. 2009. Red List
of South African Plants. *Strelitzia*
25. South African National
Biodiversity Institute, Pretoria.

Citation

Young, A.J. 2015. *Conophytum truncatum* (Thunb.) N.E.Br. subsp. *viridicatum* (N.E.Br.) S.A.Hammer. National Assessment: Red List of South African Plants version 2020.1. Accessed on 2021/10/11

 [Comment on this assessment](#)