

African Plants

[Red List statistics](#)
[methodology](#)
[Summary of recent changes](#)
[Glossary](#)
[Guidelines for EIAs](#)
[National Red List categories](#)
[Assessment](#)
Browse
[Home](#) >> [Genera: C](#) >> [Genus: Conophytum](#)

- [Genera: A](#)
- [Genera: B](#)
- [Genera: C](#)
- [Genera: D](#)
- [Genera: E](#)
- [Genera: F](#)
- [Genera: G](#)
- [Genera: H](#)
- [Genera: I](#)
- [Genera: J](#)
- [Genera: K](#)
- [Genera: L](#)
- [Genera: M](#)
- [Genera: N](#)
- [Genera: O](#)
- [Genera: P](#)
- [Genera: Q](#)
- [Genera: R](#)
- [Genera: S](#)

- [Genera: T](#)
- [Genera: U](#)
- [Genera: V](#)
- [Genera: W](#)
- [Genera: X](#)
- [Genera: Y](#)

Taxonomy

Scientific Name	Conophytum tantillum N.E.Br. subsp. lindenianum (Lavis & S.A.Hammer) S.A.Hammer
Higher Classification	Dicotyledons
Family	AIZOACEAE
Synonyms	Conophytum lindenianum Lavis & S.A.Hammer

National Status

Status and Criteria	Vulnerable D2
Assessment Date	2015/08/04
Assessor(s)	A.J. Young, D. Raimondo & L. von Staden
Justification	A range-restricted (EOO <10 km ²) habitat specialist, known from one location and potentially threatened by habitat loss to mining.

Distribution

Endemism	South African endemic
Provincial distribution	Northern Cape
Range	Namaqualand near Springbok.

Habitat and Ecology

Major system	Terrestrial
Major habitats	Namaqualand Klipkoppe Shrubland
Description	Gneiss hilltops, often inhabiting exfoliation



© A.J. Young



Search for images of **Conophytum tantillum** subsp. **lindenianum** on [iNaturalist](#)

- **General:**
- Z**

joints under rock lips
in Namaqualand
Klipkoppe Shrubland.

Threats

The population occurs near an existing mine, but is on a different geological substrate from the area that is mined. Habitat loss to infrastructure development associated with mining expansion is however a potential threat. The area is not heavily grazed and grazing does not appear to have an impact on the population.

Population

This taxon is known from a single, localized subpopulation, but it is locally common across a range of granite outcrops. Population size estimated to be more than 1 000 individuals. It quite likely also occurs in other similar habitat in the same area.

Population Stable trend

Assessment History

Taxon assessed	Status	Citation/Red List version
Criteria		
Conophytum tantillum	VU D2	2015.1
N.E.Br.		
subsp.		
lindenianum		
(Lavis & S.A.Hammer)		
S.A.Hammer		
Conophytum tantillum	Least Concern	Raimondo et al. (2009)
N.E.Br.		
subsp.		
lindenianum		
(Lavis & S.A.Hammer)		
S.A.Hammer		

Bibliography

Hammer, S. 1991. Tantalus et tantillum. Cactus and Succulent Journal (USA) 63(3):139-141.

Hammer, S. 2002. Dumpling and his wife: New view of the genus Conophytum. EAE Creative Colour, Norwich.

Hammer, S.A. 1993. The genus Conophytum: A conograph. Succulent Plant Publications, Pretoria.

Raimondo, D., von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A. and Manyama, P.A. 2009. Red List of South African Plants. *Strelitzia* 25. South African National Biodiversity Institute, Pretoria.

Citation

Young, A.J., Raimondo, D. & von Staden, L. 2015. *Conophytum tantillum* N.E.Br. subsp. *lindenianum* (Lavis & S.A.Hammer) S.A.Hammer. National Assessment: Red List of South African Plants version 2020.1. Accessed on 2021/09/28

 [Comment on this assessment](#)

(c) South African National Biodiversity Institute (**SANBI**) 2010-12.